

Amendment and Response Under 37 C.F.R. §1.116 - Expedited Examining Procedure

Page 2 of 14

Serial No.: 10/673,538

Confirmation No.: 1846

Filed: September 29, 2003

For: METHODS AND KITS FOR THE DETECTION OF ERYTHROCYTESAmendments to the Claims

This listing of claims replaces all prior versions, and listings, of claims in the above-identified application:

Listing of Claims

1. (Currently Amended) A method for detecting occult blood in a specimen comprising:
 - (a) treating the specimen with a reacting solution comprising a strong reducing agent; and
 - (b) monitoring the treated specimen for fluorescence, wherein fluorescence indicates the presence of occult blood; and wherein the treated specimen fluoresces with a spectrum from about 530 nm to about 670 nm when excited at about 480 nm.
2. (Currently Amended) A method ~~of claim 1;~~ for detecting occult blood in a specimen comprising:
 - (a) treating the specimen with a reacting solution comprising a strong reducing agent; and
 - (b) monitoring the treated specimen for fluorescence;wherein fluorescence indicates the presence of occult blood; and
wherein the strong reducing agent is sodium borohydride.
3. (Original) A method of claim 2, wherein the reacting solution is made up of approximately 0.1 percent to approximately 4 percent sodium borohydride.
4. (Original) A method of claim 2, wherein the reacting solution is made up of approximately 0.2 percent sodium borohydride.
5. (Original) A method of claim 1, wherein the reacting solution is comprised primarily of phosphate buffered saline (PBS).

Amendment and Response Under 37 C.F.R. §1.116 - Expedited Examining Procedure

Page 3 of 14

Serial No.: 10/673,538

Confirmation No.: 1846

Filed: September 29, 2003

For: METHODS AND KITS FOR THE DETECTION OF ERYTHROCYTES

6. (Previously Presented) A method of claim 1, wherein the specimen is a biological specimen selected from the group consisting of feces, urine, cerebral spinal fluid, plural cavity fluid, thoracic cavity fluid and cerebral fluid.
7. (Previously Presented) A method of claim 1, wherein the fluorescence is monitored by a fluorescent spectrometer or a fluorescent microscope.
8. (Currently Amended) A method of claim 1, wherein the ~~treated specimen fluoresces with a spectrum from about 530 to about 670 nm when excited at about 480 nm~~ strong reducing agent is sodium borohydride.
9. (Currently Amended) A method for detecting one or more erythrocytes in a specimen, wherein the method comprises:
- (a) treating the specimen with a strong reducing agent effective to enhance the fluorescence of any erythrocyte present in the specimen; and
 - (b) monitoring the fluorescence emitted by the treated specimen, wherein fluorescence of one or more erythrocytes in the treated specimen indicates the presence of erythrocytes; and wherein the treated specimen fluoresces with a spectrum from about 530 nm to about 670 nm when excited at about 480 nm.
10. (Previously Presented) A method of claim 9, wherein the specimen is a biological specimen selected from the group consisting of feces, urine, cerebral spinal fluid, plural cavity fluid, thoracic cavity fluid and cerebral fluid.
11. (Original) A method of claim 9, wherein the erythrocytes are monitored by a fluorescent microscope.

Amendment and Response Under 37 C.F.R. §1.116 - Expedited Examining Procedure

Page 4 of 14

Serial No.: 10/673,538

Confirmation No.: 1846

Filed: September 29, 2003

For: METHODS AND KITS FOR THE DETECTION OF ERYTHROCYTES

12. (Currently Amended) A method for quantifying the amount of occult blood in a specimen comprising:

- (a) exposing the specimen to a reacting solution comprising a strong reducing agent; and
- (b) monitoring the ~~treated~~ exposed specimen for fluorescence, wherein fluorescence indicates the amount of occult blood present in the specimen; and wherein the exposed specimen fluoresces with a spectrum from about 530 nm to about 670 nm when excited at about 480 nm.

13. (Previously Presented) A method of claim 1 wherein the specimen is a fecal specimen, the method further comprising, prior to step (a), purifying the fecal specimen to substantially remove all materials that will interfere with measuring the fluorescence of the fecal specimen.

14. (Currently Amended) A test kit comprising a reacting solution comprising a strong reducing agent and directions for treating a specimen with the reacting solution, wherein fluorescence of the treated specimen is indicative of the presence of occult blood in the specimen; and wherein the treated specimen fluoresces with a spectrum from about 530 nm to about 670 nm when excited at about 480 nm.

15. (Currently Amended) A ~~test kit of claim 14;~~ test kit comprising a reacting solution comprising a strong reducing agent and directions for treating a specimen with the reacting solution, wherein fluorescence of the treated specimen is indicative of the presence of occult blood in the specimen and wherein the strong reducing agent is sodium borohydride.

16. (Original) A kit of claim 15, wherein the reacting solution is made up of approximately 0.1 percent to approximately 4 percent sodium borohydride.

17. (Original) A kit of claim 15, wherein the reacting solution is made up of approximately 0.2 percent sodium borohydride.

Amendment and Response Under 37 C.F.R. §1.116 - Expedited Examining Procedure

Page 5 of 14

Serial No.: 10/673,538

Confirmation No.: 1846

Filed: September 29, 2003

For: METHODS AND KITS FOR THE DETECTION OF ERYTHROCYTES

18. (Currently Amended) A kit of claim ~~[[15]]~~ 14, wherein the reacting solution is comprised primarily of phosphate buffered saline (PBS).
19. (Previously Presented) A kit of claim 15, wherein the kit comprises containers for collecting and/or storing a specimen.
20. (Cancelled)
21. (Cancelled)
22. (Previously Presented) A kit of claim 14, wherein the kit comprises (1) directions for gathering biological samples and specimens; and (2) protocols for analyzing whether those samples and specimens contain occult blood, and if so, the amount of occult blood present in each positive sample.
23. (Original) A kit of claim 14, wherein the kit is designed to detect fecal occult blood and includes reagents and containers necessary for collecting and purifying a fecal specimen.
24. (Original) A kit of claim 14, wherein the kit comprises distinct containers for individual reagents and containers for collecting and/or storing biological samples and specimens.
25. (Original) A kit of claim 24, wherein each reagent is aliquoted in an individual container.
26. (Currently Amended) A method for comprising determining whether a subject is at risk of developing, or suffers from, a disease associated with occult blood ~~[[by]]~~ the method comprising:
- (a) treating a specimen obtained from the subject with a reacting solution comprising a strong reducing agent; and

Amendment and Response Under 37 C.F.R. §1.116 - Expedited Examining Procedure

Page 6 of 14

Serial No.: 10/673,538

Confirmation No.: 1846

Filed: September 29, 2003

For: METHODS AND KITS FOR THE DETECTION OF ERYTHROCYTES

(b) monitoring the treated specimen for fluorescence, wherein fluorescence indicates the presence of occult blood and the likelihood that the subject may develop or has developed the disease; and wherein the treated specimen fluoresces with a spectrum from about 530 nm to about 670 nm when excited at about 480 nm.

27. (Currently Amended) A method [[claim]] of claim 26, wherein the disease is gastrointestinal tumors, kidney tumors, bladder tumors, lung cancer, thoracic wall cancer, or parasite infestation and the subject is a human.

28. (Original) A method of claim 26, wherein the specimen is a biological specimen selected from the group consisting of feces, urine, cerebral spinal fluid, plural cavity fluid, thoracic cavity fluid and cerebral fluid.

29. (Previously Presented) A method for determining the extent and spatial distribution of erythrocytes trapped in cerebral tissues microvasculature comprising:

(a) treating cerebral tissue microvasculature with a reacting solution comprising a strong reducing agent; and

(b) monitoring the treated tissue for fluorescence; wherein the fluorescence indicates the extent and spatial distribution of erythrocytes trapped in cerebral tissue microvasculature.

30. (Previously Presented) A method of claim 29, wherein the vasculature is flushed with heparinized saline by cardiac perfusion to remove erythrocytes from functional post-ischemic brain microcirculation prior to treatment with the reacting solution.

31. (Currently Amended) A test kit comprising a reacting solution comprising a strong reducing agent and directions for treating cerebral tissue microvasculature with the reacting solution, wherein fluorescence of the cerebral tissue microvasculature is indicative of the extent

Amendment and Response Under 37 C.F.R. §1.116 - Expedited Examining Procedure

Page 7 of 14

Serial No.: 10/673,538

Confirmation No.: 1846

Filed: September 29, 2003

For: METHODS AND KITS FOR THE DETECTION OF ERYTHROCYTES

and spatial distribution of erythrocytes trapped in cerebral tissue microvasculature; and wherein the treated specimen fluoresces with a spectrum from about 530 nm to about 670 nm when excited at about 480 nm.

32. (Currently Amended) A method for determining whether a subject is at risk of developing, or suffers from, cerebral vascular trauma or bleeding, the method comprising:

(a) treating cerebral tissue microvasculature in situ or ex vivo with a reacting solution comprising a strong reducing agent; and

(c) (b) monitoring the treated tissue for fluorescence;

wherein the fluorescence indicates the extent and spatial distribution of erythrocytes trapped in cerebral tissue microvasculature which is indicative of the likelihood that the subject may develop or has developed the disease cerebral vascular trauma or bleeding.

33. (Currently Amended) A method of detecting the presence or past existence of erythrocytes in a specimen or sample comprising:

(a) treating the specimen or sample with a reacting solution comprising a strong reducing agent; and

(b) monitoring the treated specimen or sample for fluorescence, wherein fluorescence indicates the presence or past existence of erythrocytes in the sample or specimen; and wherein the treated specimen fluoresces with a spectrum from about 530 nm to about 670 nm when excited at about 480 nm.

34. (Currently Amended) A test kit comprising a reacting solution comprising a strong reducing agent and directions for treating a known or suspected blood sample with the reacting solution, wherein fluorescence of the treated sample is indicative of the presence of erythrocytes in the sample; and wherein the treated specimen fluoresces with a spectrum from about 530 nm to about 670 nm when excited at about 480 nm.

Amendment and Response Under 37 C.F.R. §1.116 - Expedited Examining Procedure

Page 8 of 14

Serial No.: 10/673,538

Confirmation No.: 1846

Filed: September 29, 2003

For: METHODS AND KITS FOR THE DETECTION OF ERYTHROCYTES

35. (Currently Amended) A method of claim 1, wherein the strong reducing agent is selected from the group consisting of sodium borohydride, potassium borohydride, calcium borohydride, copper borohydride, ammonium borohydride, benzyltriethylammonium borohydride, benzyl-triphenylphosphonium borohydride, bis (triphenylphosphine)copper(I) borohydride, cetyl-trimethylammonium borohydride, lithium borohydride, methytrioctylammonium borohydride, tetramethylammonium borohydride, tetrabutylammonium borohydride, tetraethylammonium borohydride, lithium aluminum hydride, diborane and 9-BBN, Dihydrogen, The Grignard Reagent, diborane, 9-BBN, dihydrogen, a grignard reagent, dialkylcopper lithium (lithium dialkylcuprate) reagents, sodium, alkyl sodium, and alkyl lithium.